RESEARCH ARTICLE

VOLCANICA

Lagged relationships between volcanic SO_2 emissions and local weather parameters in the aftermath of the 2024 Mount Lewotobi Laki-Laki eruption

Ripa Khodijah[α,γ], Suci Ramayanti[α], Andi Suhandi[α,β], Chang-Wook Lee[γ,δ], and Nanang Dwi Ardi [*α]

[α] Physics Study Program, Faculty of Mathematics and Science Education, Universitas Pendidikan Indonesia, Bandung, West Java, Indonesia.
[β] Department of Science Education, Universitas Pendidikan Indonesia, Bandung, West Java, Indonesia.
[γ] Department of Science Education, Kangwon National University, Chuncheon-si, Gangwon-do, Republic of Korea.
[δ] Department of Smart Regional Innovation, Kangwon National University, Chuncheon-si, Gangwon-do, Republic of Korea.

Submitted: 2025-09-15 | Accepted: 2026-06-16 | Published: 2026-07-02

Editor: J. Farquharson | Production: B. Hosseini, G. Seropian

Abstract

The Indonesian Centre for Volcanology and Geohazard Mitigation (PVMBG) has raised the status of Mount Lewotobi Laki-Laki to Level IV (AWAS), accompanied by massive eruptive activity with sulphur dioxide (SO_2) emissions exceeding 10,000 tons/day and an eruption column about 8,000 m high. This activity may affect local weather parameters, including surface temperature, sulphate aerosol concentration, cloud optical thickness (COT), and precipitation. To identify the influence of SO_2 emissions on these parameters, this study applies cross-correlation function (CCF) analysis using multi-sensor satellite data (OMI, AIRS, MERRA-2, MODIS, GPM). The analysis showed positive correlations with time lags: surface temperature responded after 1 day with $r = 0.50$, sulphate aerosol after 16 days with $r = 0.62$, cloud optical thickness after 21 days with $r = 0.39$, and precipitation after 23 days with $r = 0.55$. These findings demonstrate the influence of SO_2 emissions on local weather and provide valuable insights for mitigation in active regions.

Non-technical summary

Mount Lewotobi Laki-Laki in Flores, Indonesia, is currently experiencing an increase in volcanic activity and is at the highest alert level (AWAS). Satellites observed amounts of sulphur dioxide (SO_2) gas and towering ash columns. We analysed satellite data to evaluate the influence of volcanic SO_2 emissions on local weather, including surface temperature, sulphate aerosols, cloud optical thickness, and precipitation. We found that the SO_2 increase was indeed associated with these weather changes. However, the impacts occur with a lag of days to weeks. These findings suggest that volcanic eruptions can affect weather conditions in volcanically active regions.

Keywords: Mount Lewotobi Laki-Laki; SO_2 emissions; Local weather; Lag time; Satellite data.

1 Introduction

Volcanic activity, as a dynamic manifestation of Earth's internal processes, exerts a profound influence on the global atmosphere. Beyond ejecting pyroclastic materials and lava, volcanoes emit substantial quantities of gases, particularly sulphur dioxide (SO_2) [Schallock et al. 2023; Yu et al. 2023]. The injection of SO_2 into the atmosphere, especially during explosive eruptions, is one of the main precursors to the formation of sulphate (SO_4) aerosols [Marshall et al. 2020; Sellitto et al. 2024]. These aerosols play an important role in influencing weather and climate through radiative mechanisms and cloud microphysics [Toohey et al. 2021; Sahoo 2024]. Their radiative impact, especially at shortwave, can cause surface cooling [Stenchikov 2009; Lacis et al. 2013], modulate convective processes [Ceppi et al. 2017], and disrupt large-scale circulation patterns [Timmreck et al. 2021; Su et al. 2024].

While the climatic implications of volcanic sulphate aerosols are often discussed in a global context over timescales of months to years [Hermanson et al. 2020], growing evidence indicates that the most immediate atmospheric responses to volcanic SO_2 occur within the troposphere following its release. Observations across diverse volcanic systems reveal that tropospheric SO_2 exhibits strong sub-daily to short-term variability. This variability demonstrates that rapid atmospheric processing serves as a dominant short-term mechanism. Furthermore, these immediate effects operate independently of the longer term climate impact [Arellano et al. 2021; Haghighatnasab et al. 2022]. Modelling studies further demonstrate that SO_2 injected into the mid to lower troposphere is efficiently converted into sulphate aerosols through gas-phase oxidation pathways, underscoring the importance of tropospheric chemical transformation [Haghighatnasab et al. 2022; Hagen and Azevedo 2023]. The resulting enhancement in tropospheric aerosol loading substantially increases cloud droplet number concentrations, strengthening aerosol cloud interactions and modifying cloud optical properties and precipitation efficiency in the lower atmosphere [Fan et al. 2018]. These processes operate on short temporal scales and are highly sensitive to local meteorological conditions, particularly in tropical environments dominated by convection and strong moisture variability [Reid et al. 2023; Amiot et al. 2025].

Although the global climate impacts of significant volcanic eruptions have been widely studied [Stenchikov et al. 2002; Stenchikov 2009], regional atmospheric influences on the trop-

* nanang_dwiardi@upi.edu

ics remain insufficiently understood. This gap is particularly apparent in the Lesser Sunda Islands, an archipelago characterized by complex topography, monsoonal circulation, and strong ocean-atmosphere interactions [Mahrup et al. 2021; Wirasatriya et al. 2021; Simanjuntak and Lin 2022]. Disturbances from elevated SO_2 and subsequent sulphate aerosol formation can interact with local humidity, sea surface temperature, and seasonal wind patterns, affecting precipitation and convective dynamics [Li et al. 2016; Whitty et al. 2020; Sadeghi et al. 2025].

Recent studies have shown that sulphate aerosols not only increase cloud albedo and cause surface cooling, but also trigger dynamic feedbacks such as changes in latent heat release and regional circulation [Chen et al. 2024; Zheng et al. 2024; Virtanen et al. 2025]. These effects often appear with a time lag, suggesting the existence of an integrated process involving cloud microphysics and atmospheric boundary layer dynamics adjustments [Sellitto et al. 2024; Axebrink et al. 2025]. However, empirical quantification of the lag relationship between volcanic SO_2 and atmospheric variables such as surface temperature, sulphate aerosols, cloud optical thickness (COT), and precipitation is scarce, especially in tropical island environments.

The recent eruption of Mount Lewotobi Laki-Laki (November 2024) provides an ideal natural study site to analyse this phenomenon. On 3 November, 2024, the Indonesian Centre for Volcanology and Geohazard Mitigation (PVMBG) raised the alert status to Level IV (AWAS) after an eruption that spewed an ash column 8 km high and released more than 10,000 tons/day of SO_2 [Ministry of Energy and Mineral Resources 2024a; b]. This high SO_2 flux has the potential to disrupt regional weather conditions, triggering a two-phase thermal response: an initial warming phase through radiation absorption followed by a delayed cooling phase due to increased cloud cover and altering precipitation patterns through microphysical mechanisms and atmospheric dynamics [Malavelle et al. 2017; Marshall et al. 2022; Paik et al. 2023; Lin et al. 2025].

Most previous studies have focused on global climate modelling. Few studies have utilized a statistical framework to detect the lagged atmospheric response of volcanic SO_2 disturbances. Findings from an urban air pollution study show that SO_2 has significant delayed correlations with surface temperature (positive at +3, +5, +13, and +16 days) and with precipitation (negative after +1 day) [Liu et al. 2025]. Although providing valuable insights, this approach has had limited application to volcanic studies on tropical islands.

To address this gap, this study integrates multi-satellite observations with cross-correlation function (CCF) analysis to quantify the temporal and spatial linkages between volcanic SO_2 and key weather variables in the Lesser Sundas. In particular, we detect lag-dependent sequential associations between SO_2, sulphate aerosols, COT, surface temperature, and precipitation, and visualize the spatial heterogeneity of atmospheric response following volcanic SO_2 emissions. This integrative framework provides new insights into the short-term atmospheric impacts of volcanic SO_2, improves regional weather understanding, and supports disaster risk mitigation strategies.

2 Data and Methodology

This study employs an integrated framework of satellite-based observations and statistical analysis to assess the atmospheric response to volcanic SO_2 emissions from Mount Lewotobi Laki-Laki. Daily datasets from OMI, AIRIS, MERRA-2, MODIS, and GPM (all defined below) underwent preprocessing through noise filtering and temporal resampling to obtain consistent daily resolution. We employed CCF analysis to quantify the lag-dependent associations between SO_2, surface temperature, sulphate aerosol, COT, and precipitation. This approach enables systematic detection of sequential atmospheric responses, while providing new insights into the timing and intensity of aerosol, cloud, and precipitation interactions following volcanic emissions. The methodological workflow is summarized in Figure 1, which highlights the stages from raw satellite data acquisition to final interpretation in the temporal and spatial domains.




Figure 1: Flowchart of the methodology illustrating the analytical framework of the study, starting with the acquisition of satellite data (SO_2, temperature, aerosols, cloud optical thickness (COT), and precipitation), followed by the process of data filtering (noise removal) and temporal alignment, cross-correlation function (CCF) analysis on weather variables, and ending with temporal interpretation and spatial visualization.

2.1 Data acquisition

This study examines the atmospheric impact of post-eruption volcanic SO_2 emissions of Mount Lewotobi Laki-Laki by analysing datasets extracted from various satellite instruments via the Giovanni platform*. The total SO_2 column, which serves as the independent variable, was obtained from the Ozone Monitoring Instrument (OMI) operating on the Aura satellite†. The dependent variable is the surface temperature of the lower troposphere, which was obtained from the Atmospheric Infrared Sounder (AIRS) on the Aqua satellite‡. Sulphate aerosol concentrations were derived via the MERRA-2 reanalysis product§. Cloud optical thickness (COT) data were retrieved from the Moderate Resolution Imaging Spectroradiometer (MODIS) instrument on the Aqua and Terra satellites¶. Daily precipitation data were sourced from the Global Precipitation Measurement (GPM) mission a collaboration between NASA and JAXA‖. The analysis period spans 1 October to 31 December, 2024, allowing for a comprehensive assessment of weather conditions before, during, and after the eruption.

2.2 Study area

This study focuses on the Lesser Sunda Islands, an active volcanic archipelago in eastern Indonesia spanning a total land area of approximately 75,000 km^2. The region includes the main islands of Flores, Sumba, Timor, Savu, and Lembata, along with their surrounding marine environments. It is bordered by the Banda Sea to the north, the Savu Sea to the south, and the Flores Sea and Timor Sea to the east and west [Mahrup et al. 2021]. A spatial overview of the study site, illustrating the position of Mount Lewotobi Laki-Laki and the geographic setting of the Lesser Sunda Islands, is presented in Figure 2.

The Lesser Sunda region experiences a tropical monsoon climate driven by the Asia–Australia monsoon system. During the wet season (October to April), north-westerly winds transport moisture from the Indian and Pacific Oceans. In contrast, dry easterlies from mainland Australia dominate the dry season (May to September) [Trismidianto et al. 2024]. Mean annual temperatures range between 26 and 28 °C, with substantial precipitation variability. Recent climatological analyses indicate a decline in precipitation linked to rising sea surface temperatures [Mahrup et al. 2021]. Furthermore, previous studies in Indonesia have demonstrated that large-scale atmospheric phenomena such as tropical cyclones can significantly alter local meteorological parameters, including temperature, pressure, and humidity [Nugroho et al. 2019], underscoring the sensitivity of the regional climate system to external forcings.

Topographically, the region consists of rugged volcanic mountains, steep coastal slopes, and limited lowlands. Mount Lewotobi Laki-Laki (1,584 m a.s.l.) and Mount Lewotobi Perempuan (1,703 m a.s.l.) are paired active volcanoes in East Flores [Global Volcanism Program 2024], forming part




Figure 2: [A] Geographical location of the Lesser Sunda Islands, including the mainland and surrounding seas; [B] Position of Mount Lewotobi Laki-Laki and Mount Lewotobi Perempuan.

of the Sunda Arc, a subduction-related volcanic chain generated by the convergence of the Indo-Australian and Eurasian plates [Hall 2012]. Oceanographically, the region is subject to seasonal upwelling and the Indonesian Throughflow, which regulates local sea surface temperatures and can affect post-eruption aerosol dispersion [Wirasatriya et al. 2021]. The interactions between volcanic activity, monsoon circulation, and the complex topography of the islands make this an ideal region to evaluate the influence of volcanic SO_2 emissions on local weather dynamics through lag correlation analysis using the CCF.

2.3 Cross-correlation function (CCF) analysis

The primary analysis in this study uses the CCF to evaluate the temporal relationship between volcanic SO_2 emissions and local weather variables, including surface temperature, sulphate aerosol, COT, and precipitation. The CCF calculates the Pearson correlation coefficient r between the SO_2 time series and each local weather parameter at various lags τ, thereby identifying the strength and direction of the delayed relationship. The r value at each lag was interpreted using Papageorgiou [2022] classification: $r > 0.9$ (almost perfect), $0.7 < r \leq 0.9$ (very strong), $0.5 < r \leq 0.7$ (strong), $0.3 < r \leq 0.5$ (moderate), $0.1 < r \leq 0.3$ (weak), and $r < 0.1$ (very weak). Mathematically, the CCF for two discrete time series, $x(t)$ and $y(t)$, at lag τ is defined as:

$$\mathrm{CCF}(\tau) = \frac{\sum_{t=1}^{N-1} [(x(t) - \bar{x})(y(t-\tau) - \bar{y})]}{\sqrt{\sum_{t=1}^{N-1} (x(t) - \bar{x})^2 \sum_{t=1}^{N-1} (y(t-\tau) - \bar{y})^2}}, \quad (1)$$

where $\bar{x}$ and $\bar{y}$ are the mean values and N is the number of observations [Hoffmann et al. 2024]. The lag τ with the highest correlation coefficient r indicates the time lag of the

*https://giovanni.gsfc.nasa.gov/
†https://aura.gsfc.nasa.gov/omi.html
‡https://airs.jpl.nasa.gov/
§https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/
¶https://modis.gsfc.nasa.gov/
‖https://gpm.nasa.gov/

atmospheric response to peak SO_2 emissions. A positive correlation reflects an increase in weather variables concurrent with rising SO_2 emissions, while a negative correlation signifies an inverse response.

3 Results

3.1 Daily observations of SO_2 emissions and weather parameters

Daily observations of the atmospheric response were analysed by representing daily values spatially averaged over a fixed regional domain (115–125°E, 8–11°S). This specific domain was selected to encompass Mount Lewotobi and its dominant downwind transport pathway under prevailing monsoonal circulation, ensuring that the non-stationary nature of the SO_2 plume was captured within the study area. Furthermore, the analysis was systematically segmented into three temporal phases to provide a consistent atmospheric framework: the pre-eruption period (1 October to 2 November), the eruption phase (3 November to 3 December), and the post-eruption period (4–31 December). During the eruptive phase, SO_2 concentrations exhibited a sharp increase, peaking at approximately 50.62 kt (Figure 3A). This abrupt rise triggered a sequence of atmospheric responses. During the eruption, surface temperatures initially climbed to about 27.6 °C (Figure 3B), likely driven by direct radiative heating from volcanic gas emissions and particle interactions in the atmosphere. Following the emission peak, temperatures did not immediately return to baseline but gradually declined to approximately 25.0 °C in the post-eruption period (Figure 3B). This pattern reflects a two-stage thermal response: initial warming followed by delayed cooling, mediated by the formation of sulphate aerosols.

Sulphate aerosol concentrations began to increase several days after the SO_2 peak, and reached a maximum of 1.7×10^{-5} kg m^{-2} (Figure 3C). This delay reflects the time required for the chemical reaction of SO_2 oxidation in the atmosphere, and confirms that sulphate aerosols have a longer residence time in the atmosphere. These aerosols also affected cloud properties, as evidenced by a substantial increase in COT, which peaked at 22.1 (Figure 3D), substantially higher than the pre-eruption daily range of 5-–10. This increase is in line with the Twomey effect, where the number of small cloud droplets increases as sulphate aerosols act as cloud condensation nuclei (CCN), resulting in thicker and more reflective clouds. Even after SO_2 emissions decreased, COT values remained high for several days before returning to normal, confirming the role of persistent aerosols in modifying cloud structure.

Meanwhile, precipitation exhibited an opposite trend. Before the eruption, daily precipitation was relatively low, averaging 0–4 mm/day, with a brief spike of approximately 28 mm/day in late October, likely marking the onset of the Lesser Sunda rainy season [Mahrup et al. 2021]. During the eruptive phase, precipitation declined sharply to 0–5 mm/day and occasionally dropped to near zero, despite otherwise favorable atmospheric conditions for cloud formation. This pattern suggests a disruption of raindrop coalescence caused by elevated concentrations of sulphate aerosols acting as CCN. Following the decline of SO_2 emissions, precipitation increased substantially, reaching a peak of ~38 mm/day in early December (Figure 3E). Such a sequence reflects a lagged hydrological response to volcanic SO_2 emissions, in which aerosol injection temporarily suppresses precipitation before intensification resumes as atmospheric conditions stabilize. The temporal evolution of this response is systematically illustrated in Figure 3A–3E, highlighting the interconnections between SO_2 dynamics and key weather variables.

3.2 Responses: cross-correlation analysis

CCF analysis between volcanic SO_2 emissions and atmospheric parameters reveals distinct temporal response patterns, reflecting the sequential nature of atmospheric processes triggered by eruptive activity. To establish a temporal baseline, the analysis utilizes the peak SO_2 emission observed on 10 November 2024 (as shown in Figure 3A) as the reference point ($\tau = 0$). In this context, the lag values represent the duration required for the eruptive peak to generate a maximum response or peak pattern in each meteorological parameter. The fastest response appears in surface temperature (Figure 4A), which reaches a maximum correlation of $r = 0.51$ at a 1 day lag (11 November). This finding indicates that SO_2 immediately affects the surface radiation balance, either through direct absorption of solar radiation or through initial interactions with cloud cover. After peaking, the correlation declines sharply and enters negative values between lags of 15–35 days, suggesting that the short lived radiative effects of SO_2 are given way to by secondary processes, such as the formation of sulphate aerosols from atmospheric SO_2 oxidation, which can increase cloud albedo and reduce incoming solar radiation at the surface.

A subsequent response emerges in sulphate (SO_4) aerosols (Figure 4B), reaching a peak correlation of $r = 0.62$ at a 16-day lag (26 November). This delay aligns with the atmospheric conversion of SO_2 into secondary sulphate particles through oxidation processes. Once generated, these aerosols influence cloud microphysics, as evidenced by a peak in COT at day 21 (1 December), showing a correlation of $r = 0.39$ (Figure 4C). The rise in COT reflects an enhanced cloud droplet number and greater optical density, driven by the role of aerosols as CCNs.

Precipitation exhibits the most delayed response (Figure 4D), reaching a peak correlation of $r = 0.55$ at a 23-day lag which closely matches the significant rainfall intensification observed on 3 December (Figure 3E). This behaviour aligns with the final stage of the aerosol-cloud-precipitation feedback cycle, in which modifications to cloud microphysics ultimately enhance precipitation efficiency. The progression of peaks spanning rapid radiative forcing, chemical transformation to aerosols, cloud adjustment, and precipitation modulation depicts a complete atmospheric pathway initiated by volcanic SO_2 emissions. This sequential relationship is clearly illustrated in Figure 4A–4D.




Figure 3: Daily variations of SO_2 emissions (kilotons), and local weather parameters including surface temperature (°C), SO_4 mass ($\times 10^{-5}$ kg m^{-2}), cloud optical thickness (COT), and precipitation (mm/day). The gray area marks the eruption period (3 November to 3 December, 2024), which could potentially affect the observed atmospheric dynamics.

4 Discussion

Daily observations of surface temperature indicate that although the Lesser Sunda Islands experience dynamically varying daily temperature fluctuations, the peak of 27.59 °C recorded in November represents a pronounced thermal anomaly that spatially coincides with the eruptive activity of Mount Lewotobi (Figure 3B). This value is particularly significant as it exceeds the regional climatological baseline for the early wet season transition, which typically remains below 27.5 °C [Simanjuntak and Lin 2022]. Physically, this elevated temperature suggests an additional thermal energy input via the volcanic blanket effect, where volcanic gases and ash trap outgoing terrestrial longwave radiation [Sellitto et al. 2024]. The robustness of this anomaly is further supported by the CCF analysis, which reveals a very short lag of only 1 day between SO_2 emissions and the observed temperature increase. Evidence of this localized energy enhancement is clearly illustrated in the November transect profile (Figure 5E), where the most pronounced temperature increase is concentrated specifically from the crater extending up to 3 km eastward, corresponding to the location of the eruption source.

However, as the eruption progresses, this warming phase is temporary. The subsequent gradual cooling toward ~25 °C during the later stages of the eruption period aligns with the peak accumulation of sulphate aerosols (Figure 3C). This ~25 °C stabilization is consistent with the climatological baseline for the Lesser Sunda Islands region during this transition, which typically ranges between 25 °C and 27.5 °C [Simanjuntak and Lin 2022]. This transition is driven by atmospheric chemical transformations where the initial thermal phase decays and promotes the formation of sulphate aerosols, reaching their maximum correlation at a 16-day lag ($r = 0.62$). This pattern reflects oxidation processes involving hydroxyl radicals (OH) under ultraviolet radiation, combined with the in-




Figure 4: Cross-correlation analysis between volcanic SO_2 emissions and four local weather parameters: [A] surface temperature, [B] SO_4 aerosol, [C] cloud optical thickness (COT), and [D] precipitation, during the Mount Lewotobi Laki-Laki observation period in 2024. The x-axis shows the time lag (in days), where a positive lag indicates that changes in SO_2 precede changes in the associated atmospheric parameters. Each panel displays the maximum correlation coefficient (r) along with its optimal lag, marked by the dashed vertical line and the highlighted dot. The gray shaded area represents the ±95% confidence interval, where correlation values within this range are statistically insignificant.

fluence of ambient atmospheric humidity, consistent with the estimated e-folding lifetime of volcanic SO_2 of approximately 13–17 days [Cai et al. 2022]. In response to the inherently non-stationary nature of volcanic plumes, spatial analysis (Figure 5H) demonstrates that these aerosols do not remain confined above the crater but are transported by prevailing winds moving from east to west. The resulting accumulation of sulphate aerosols subsequently acts as a catalyst for cloud microphysical modification, leading to an increase in COT with a 21-day lag (r = 0.39). This elevation in COT, despite the absence of immediate precipitation, indicates the formation of an optically dense yet meteorologically stable cloud layer, a phenomenon driven by the high concentration of volcanic aerosols. The conceptual framework for these microphysical modifications is further detailed in Figure 6.

From a microphysical perspective, the massive injection of fine aerosols (<1 µm) intensifies competition for available water vapour, producing cloud droplets with high number con-




Figure 5: (Caption next page.)

Figure 5: (Previous page.) Spatiotemporal synthesis of atmospheric responses. The top panels of each sub-figure represent the monthly average spatial distribution, featuring a red triangle as the volcano location and a dotted line indicating the transect path. The bottom panels display the corresponding monthly average longitudinal transects extending from the volcano location (0 km) to across the upwind and downwind sectors for October (baseline), November (syn-eruption), and December (post-eruption). The observed changes in the November and December phases correspond to the identified peak lags of SO_2 (1-day), SO_4 mass (16-day), cloud optical thickness (21-day), and precipitation (23-day), respectively.




Figure 6: Conceptual diagram of cloud microphysical modifications. The illustration depicts the transition from SO_2 to sulphate aerosols acting as cloud condensation nuclei (CCN), promoting droplet competition and suppressed coalescence (Twomey effect), which increases cloud optical thickness (COT) without immediately producing precipitation.

centrations but very small diameters, thereby inhibiting efficient coalescence processes. This mechanism explains the anomaly observed in Figure 3, where despite the region having already passed the natural monsoon onset during the final week of October (rainfall peak of ~28 mm/day), precipitation throughout November was markedly suppressed, approaching near zero values. Under these conditions, water mass remained effectively suspended within the atmosphere, while air masses were advected from the crater source region (0 km) toward the downstream sector (0–33 km), as detailed in the conceptual framework in Figure 6.

The hydrological response ultimately reached a saturation point in December, when the accumulated atmospheric water mass was released as extreme precipitation reaching 38.29 mm/day. The observed 23-day lag represents the physical timescale required for the air mass to travel across the east to west transport pathway while undergoing progressive microphysical maturation. Unlike the sporadic rainfall events observed in October, the December precipitation burst was both intense and spatially displaced toward the 0–33 km sector (Figure 5O), reflecting the downwind transport pathways established during the peak SO_2 loading in November. This phenomenon provides clear evidence of an aerosol indirect effect, whereby interactions between volcanic emissions and atmospheric transport dynamics fundamentally altered the normal seasonal rainfall pattern, ultimately producing an extreme hydrometeorological response in the downwind region [Wong et al. 2025]. Similar short-term meteorological anomalies have been documented following the 2022 Hunga Tonga-Hunga Haʻapai eruption, where a maturation period of several weeks was required for atmospheric cooling and cloud formation to culminate in record-breaking rainfall [Wong and Yim 2022]. Furthermore, our observed 23-day lag closely matches historical records where distal regions, such as Hong Kong, recorded precipitation delays of 23 to 35 days following the 1982 El Chichón and 2008 Chaitén eruptions [Yim 2005; Wong et al. 2023]. While the Tongan event reached stratospheric heights and involved massive water vapour, the Mount Lewotobi Laki-Laki case demonstrates that tropospheric sulphur loading can trigger comparable physical maturation timescales, leading to similar downwind precipitation bursts. The synthesized relationship between these spatiotemporal variables is illustrated in the schematic and transect profiles in Figure 5.

5 Conclusions

The November 2024 eruption of Mount Lewotobi Laki-Laki triggered a temporally and spatially phased series of atmospheric responses. The initial phase exhibited a brief warming of surface temperature (~27.6 °C) with a maximum correlation of $r = 0.51$ at a lag of 1 day, triggered by the volcanic blanket effect, in which volcanic gases and ash absorb and trap Earth's radiation, resulting in localized warming. This effect is temporary, followed by gradual cooling (~25 °C) as atmospheric reflectivity increases due to sulphate aerosol formation, consistent with the volcanic umbrella effect. SO_2 reaches its highest correlation with sulphate aerosols, $r = 0.62$ at a lag of 16 days, reflecting gradual chemical conversion through atmospheric oxidation. Mature aerosols act as cloud condensation nuclei (CCN), enhancing cloud optical thickness (COT) with a peak correlation of $r = 0.39$ at a 21-day lag, making clouds more reflective and durable. The hydrological response appears to be the slowest, characterized by a two-phase pattern: initially suppressing precipitation during the degassing phase, then increasing to ~38 mm/day with a peak correlation of $r = 0.55$ at 23 days lag. Spatial analysis, conducted via longitudinal transects, reveals that these impacts are not stationary; rather, they follow a dominant east-to-west transport mechanism. While thermal anomalies were localized at the crater and extending up to 3 km eastward, the subsequent aerosol maturation and microphysical modifications triggered a "downwind shift" in precipitation following November's transport pathways. This process culminated in extreme rainfall in the distal downstream sectors (0–30 km) in December, rather than the immediate volcanic vicinity.

These findings confirm that aerosols modulate local weather through a sequence of rapid warming, atmospheric cooling, cloud modification, and delayed precipitation. Monitoring SO_2 and sulphate aerosols is therefore critical for supporting early warnings of extreme precipitation following eruptive surges. However, this study is limited by the lack of atmospheric vertical profile data and in situ validation, which prevents a complete understanding of altitude dynamics and cloud microphysical interactions. Consequently, further research should integrate WRF-Chem-based atmospheric modelling, multi-sensor satellite observations, and field measurements. The integration of these approaches, combined with longer observation periods, will also allow for advanced statistical verification such as the Granger causality test. Furthermore, future studies could explore the comparative impacts of subaerial and submarine eruptions, specifically how large-scale water vapour emissions in submarine events might influence the time-lag of sulphate formation and weather responses. This will enable aerosol, cloud, and precipitation feedbacks to be examined more comprehensively and improve the accuracy of volcanic weather impact predictions.

Author Contributions

Conceptualization: R.K., and S.R.; Methodology: R.K., and N.D.A.; Software: R.K.; Formal analysis: R.K., and C.W.L.; Investigation: R.K. and S.R.; Data curation: R.K. and A.S.; Validation: A.S., N.D.A., and C.W.L.; Resources: A.S. and N.D.A.; Writing original draft: R.K.; Writing review & editing: C.W.L., and N.D.A.; Visualization: R.K., and S.R.; Supervision: A.S. and N.D.A.; Project administration: A.S.; Funding acquisition: N.D.A, and A.S.

Acknowledgements

The authors express their sincere gratitude to the NASA Earth Science Data Systems (ESDS) for providing open-access satellite observations through the Giovanni platform, particularly from missions such as OMI, MODIS, MERRA-2, AIRS and GPM, which were essential for the atmospheric analysis in this study. Appreciation is also extended to the Centre for Volcanology and Geohazard Mitigation (PVMBG) for publicly available eruption reports that supported contextual interpretation of the volcanic activity.

Data Availability

All raw satellite data utilized in this study (SO_2, sulphate aerosol concentrations, surface temperature, cloud optical thickness (COT), and precipitation) are publicly available through the NASA Goddard Earth Sciences Data and Information Services Center (GES DISC), accessed via the Giovanni platform (https://giovanni.gsfc.nasa.gov/giovanni/). The processed and derived datasets generated in this research are available from the corresponding author upon reasonable request.

Copyright Notice

© The Author(s) 2026. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

Amiot, C. G., T. J. Lang, S. C. van den Heever, R. A. Ferrare, O. O. Sy, L. D. Carey, S. A. Christopher, J. R. Mecikalski, S. W. Freeman, G. A. Sokolowsky, C. A. Hostetler, and S. Tanelli (2025). "Observed impacts of aerosol concentration on maritime tropical convection within constrained environments using airborne radiometer, radar, lidar, and dropsondes". In: *Atmospheric Chemistry and Physics* 25(19), pages 12335–12355. DOI: 10.5194/acp-25-12335-2025.

Arellano, S., B. Galle, F. Apaza, G. Avard, C. Barrington, N. Bobrowski, C. Bucarey, V. Burbano, M. Burton, Z. Chacón, G. Chigna, C. J. Clarito, V. Conde, F. Costa, M. De Moor, H. Delgado-Granados, A. Di Muro, D. Fernandez, G. Garzón, H. Gunawan, N. Haerani, T. H. Hansteen, S. Hidalgo, S. Inguaggiato, M. Johansson, C. Kern, M. Kihlman, P. Kowalski, P. Masias, F. Montalvo, J. Möller, U. Platt, C. Rivera, A. Saballos, G. Salerno, B. Taisne, F. Vásconez, G. Velásquez, F. Vita, and M. Yalire (2021). "Synoptic analysis of a decade of daily measurements of SO_2 emission in the troposphere from volcanoes of the global ground-based Network for Observation of Volcanic and Atmospheric Change". In: *Earth System*

Science Data 13(3), pages 1167–1188. DOI: 10.5194/essd-13-1167-2021.

Axebrink, E., M. K. Sporre, and J. Friberg (2025). "Impact of SO_2 injection profiles on simulated volcanic forcing for the 2009 Sarychev eruptions – investigating the importance of using high-vertical-resolution methods when compiling SO_2 data". In: *Atmospheric Chemistry and Physics* 25(4), pages 2047–2059. DOI: 10.5194/acp-25-2047-2025.

Cai, Z., S. Griessbach, and L. Hoffmann (2022). "Improved estimation of volcanic SO_2 injections from satellite retrievals and Lagrangian transport simulations: the 2019 Raikoke eruption". In: *Atmospheric Chemistry and Physics* 22(10), pages 6787–6809. DOI: 10.5194/acp-22-6787-2022.

Ceppi, P., F. Brient, M. D. Zelinka, and D. L. Hartmann (2017). "Cloud feedback mechanisms and their representation in global climate models". In: *WIREs Climate Change* 8(4). DOI: 10.1002/wcc.465.

Chen, Y., J. Haywood, Y. Wang, F. Malavelle, G. Jordan, A. Peace, D. G. Partridge, N. Cho, L. Oreopoulos, D. Grosvenor, P. Field, R. P. Allan, and U. Lohmann (2024). "Substantial cooling effect from aerosol-induced increase in tropical marine cloud cover". In: *Nature Geoscience* 17(5), pages 404–410. DOI: 10.1038/s41561-024-01427-z.

Fan, C., M. Ding, P. Wu, and Y. Fan (2018). *The Relationship between Precipitation and Aerosol: Evidence from Satellite Observation*. DOI: 10.48550/ARXIV.1812.02036.

Global Volcanism Program (2024). "Lewotobi (264180)". In: *[Database] Volcanoes of the World (v. 5.2.3; 20 Sep 2024).* Distributed by Smithsonian Institution, compiled by Venzke, E. DOI: 10.5479/si.GVP.VOTW5-2024.5.2.

Hagen, M. and A. Azevedo (2023). "Influence of Volcanic Activity on Weather and Climate Changes". In: *Atmospheric and Climate Sciences* 13(02), pages 138–158. DOI: 10.4236/acs.2023.132009.

Haghighatnasab, M., J. Kretzschmar, K. Block, and J. Quaas (2022). "Impact of Holuhraun volcano aerosols on clouds in cloud-system-resolving simulations". In: *Atmospheric Chemistry and Physics* 22(13), pages 8457–8472. DOI: 10.5194/acp-22-8457-2022.

Hall, R. (2012). "Late Jurassic–Cenozoic reconstructions of the Indonesian region and the Indian Ocean". In: *Tectonophysics* 570-571, pages 1–41. DOI: 10.1016/j.tecto.2012.04.021.

Hermanson, L., R. Bilbao, N. Dunstone, M. Ménégoz, P. Ortega, H. Pohlmann, J. I. Robson, D. M. Smith, G. Strand, C. Timmreck, S. Yeager, and G. Danabasoglu (2020). "Robust Multiyear Climate Impacts of Volcanic Eruptions in Decadal Prediction Systems". In: *Journal of Geophysical Research: Atmospheres* 125(9). DOI: 10.1029/2019jd031739.

Hoffmann, L., L. Gilardi, M.-T. Schmitz, T. Erbertseder, M. Bittner, S. Wüst, M. Schmid, and J. Rittweger (2024). "Investigating the spatiotemporal associations between meteorological conditions and air pollution in the federal state Baden-Württemberg (Germany)". In: *Scientific Reports* 14(1). DOI: 10.1038/s41598-024-56513-4.

Lacis, A. A., J. E. Hansen, G. l. Russell, V. Oinas, and J. Jonas (2013). "The role of long-lived greenhouse gases as principal LW control knob that governs the global surface temperature for past and future climate change". In: *Tellus B: Chemical and Physical Meteorology* 65(1), page 19734. DOI: 10.3402/tellusb.v65i0.19734.

Li, Z., W. K.-M. Lau, V. Ramanathan, G. Wu, Y. Ding, M. G. Manoj, J. Liu, Y. Qian, J. Li, T. Zhou, J. Fan, D. Rosenfeld, Y. Ming, Y. Wang, J. Huang, B. Wang, X. Xu, S.-S. Lee, M. Cribb, F. Zhang, X. Yang, C. Zhao, T. Takemura, K. Wang, X. Xia, Y. Yin, H. Zhang, J. Guo, P. M. Zhai, N. Sugimoto, S. S. Babu, and G. P. Brasseur (2016). "Aerosol and monsoon climate interactions over Asia". In: *Reviews of Geophysics* 54(4), pages 866–929. DOI: 10.1002/2015rg000500.

Lin, C., J. Z. Yu, E. Lee, P.-W. Chan, J. W.-Y. Ng, Y.-W. Chan, T. Zhang, Y. Chen, H. Chen, Z. Zhang, S. H.-K. Chui, J. C. H. Fung, W. Liu, and A. K. H. Lau (2025). "Mysterious air pollution in south China linked to volcanic emissions from the Philippines". In: *Communications Earth & Environment* 6(1). DOI: 10.1038/s43247-025-02073-y.

Liu, X., J. Jeon, A. R. Nacht, and P. S. Larson (2025). "Lag Associations of Precipitation and Temperature with Seven Types of Ambient Air Pollution Concentrations in the Midwestern United States: A Time Series Analysis". In: DOI: 10.20944/preprints202503.0438.v1.

Mahrup, M. Ma'shum, and Fahrudin (2021). "Climate change in The Lesser Sunda Islands: The harsh region in the maritim continent of Indonesia". In: *IOP Conference Series: Earth and Environmental Science* 824(1), page 012115. DOI: 10.1088/1755-1315/824/1/012115.

Malavelle, F. F., J. M. Haywood, A. Jones, A. Gettelman, L. Clarisse, S. Bauduin, R. P. Allan, I. H. H. Karset, J. E. Kristjánsson, L. Oreopoulos, N. Cho, D. Lee, N. Bellouin, O. Boucher, D. P. Grosvenor, K. S. Carslaw, S. Dhomse, G. W. Mann, A. Schmidt, H. Coe, M. E. Hartley, M. Dalvi, A. A. Hill, B. T. Johnson, C. E. Johnson, J. R. Knight, F. M. O'Connor, D. G. Partridge, P. Stier, G. Myhre, S. Platnick, G. L. Stephens, H. Takahashi, and T. Thordarson (2017). "Strong constraints on aerosol–cloud interactions from volcanic eruptions". In: *Nature* 546(7659), pages 485–491. DOI: 10.1038/nature22974.

Marshall, L. R., E. C. Maters, A. Schmidt, C. Timmreck, A. Robock, and M. Toohey (2022). "Volcanic effects on climate: recent advances and future avenues". In: *Bulletin of Volcanology* 84(5). DOI: 10.1007/s00445-022-01559-3.

Marshall, L. R., C. J. Smith, P. M. Forster, T. J. Aubry, T. Andrews, and A. Schmidt (2020). "Large Variations in Volcanic Aerosol Forcing Efficiency Due to Eruption Source Parameters and Rapid Adjustments". In: *Geophysical Research Letters* 47(19). DOI: 10.1029/2020gl090241.

Ministry of Energy and Mineral Resources (2024a). *Aktivitas Vulkanik Meningkat, PVMBG Naikkan Status G. Lewotobi Laki-laki Naik Jadi "AWAS." Published 04/11/2024.* URL: https://www.esdm.go.id/id/media-center/arsip-berita/aktivitas-vulkanik-meningkat-pvmbg-naikkan-status-g-lewotobi-laki-laki-naik-jadi-awas?fbclid=IwY2xjawGVGbxleHRuA2FlbQIxMAABHTCZeDBpIpHCPLy-qwqC-9TsXknMdFqQi5PCUrutLIbXdsm9IOO5N7xgpg_aem_dLMg2xZ80VImL0bK6.

– (2024b). *Badan Geologi: Erupsi G. Lewotobi Laki-Laki Keluarkan Gas SO_2 Lebih dari 10.000 Ton per Hari. Published 04/11/2024*. URL: https://www.esdm.go.id/en/media-center/news-archives/badan-geologi-erupsi-g-lewotobi-laki-laki-keluarkan-gas-so2-lebih-dari-10000-ton-per-hari.

Nugroho, H. S., C. P. Asmoro, A. R. Achmad, N. D. Ardi, A. F. C. Wijaya, J. A. Utama, and Y. R. Tayubi (2019). "How tropical cyclones affected temperature, pressure, and humidity in Universitas Pendidikan Indonesia". In: *Journal of Physics: Conference Series* 1280(2), page 022077. DOI: 10.1088/1742-6596/1280/2/022077.

Paik, S., S.-K. Min, S.-W. Son, E.-P. Lim, S. McGregor, S.-I. An, J.-S. Kug, and S.-W. Yeh (2023). "Impact of volcanic eruptions on extratropical atmospheric circulations: review, revisit and future directions". In: *Environmental Research Letters* 18(6), page 063003. DOI: 10.1088/1748-9326/acd5e6.

Papageorgiou, S. N. (2022). "On correlation coefficients and their interpretation". In: *Journal of Orthodontics* 49(3), pages 359–361. DOI: 10.1177/14653125221076142.

Reid, J. S., H. B. Maring, G. T. Narisma, S. van den Heever, L. Di Girolamo, R. Ferrare, P. Lawson, G. G. Mace, J. B. Simpas, S. Tanelli, L. Ziemba, B. van Diedenhoven, R. Bruintjes, A. Bucholtz, B. Cairns, M. O. Cambaliza, G. Chen, G. S. Diskin, J. H. Flynn, C. A. Hostetler, R. E. Holz, T. J. Lang, K. S. Schmidt, G. Smith, A. Sorooshian, E. J. Thompson, K. L. Thornhill, C. Trepte, J. Wang, S. Woods, S. Yoon, M. Alexandrov, S. Alvarez, C. G. Amiot, J. R. Bennett, M. Brooks, S. P. Burton, E. Cayanan, H. Chen, A. Collow, E. Crosbie, A. DaSilva, J. P. DiGangi, D. D. Flagg, S. W. Freeman, D. Fu, E. Fukada, M. R. A. Hilario, Y. Hong, S. M. Hristova-Veleva, R. Kuehn, R. S. Kowch, G. R. Leung, J. Loveridge, K. Meyer, R. M. Miller, M. J. Montes, J. N. Moum, A. Nenes, S. W. Nesbitt, M. Norgren, E. P. Nowottnick, R. M. Rauber, E. A. Reid, S. Rutledge, J. S. Schlosser, T. T. Sekiyama, M. A. Shook, G. A. Sokolowsky, S. A. Stamnes, T. Y. Tanaka, A. Wasilewski, P. Xian, Q. Xiao, Z. Xu, and J. Zavaleta (2023). "The Coupling Between Tropical Meteorology, Aerosol Lifecycle, Convection, and Radiation during the Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP2Ex)". In: *Bulletin of the American Meteorological Society* 104(6), E1179–E1205. DOI: 10.1175/bams-d-21-0285.1.

Sadeghi, B., A. Crawford, T. Chai, M. Cohen, J. Sieglaff, M. Pavolonis, H. C. Kim, and G. Morris (2025). "Improving Volcanic SO_2 Cloud Modeling Through Data Fusion and Trajectory Analysis: A Case Study of the 2022 Hunga Tonga Eruption". In: *Journal of Geophysical Research: Atmospheres* 130(4). DOI: 10.1029/2024jd042421.

Sahoo, S. S. (2024). "Volcanic eruptions and climate: A complex interplay". In: *Journal of Geosciences Insights* 2(1), pages 25–28. DOI: 10.61577/jgi.2024.100005.

Schallock, J., C. Brühl, C. Bingen, M. Höpfner, L. Rieger, and J. Lelieveld (2023). "Reconstructing volcanic radiative forcing since 1990, using a comprehensive emission inventory and spatially resolved sulfur injections from satellite data in a chemistry-climate model". In: *Atmospheric Chemistry and Physics* 23(2), pages 1169–1207. DOI: 10.5194/acp-23-1169-2023.

Sellitto, P., R. Siddans, R. Belhadji, E. Carboni, B. Legras, A. Podglajen, C. Duchamp, and B. Kerridge (2024). "Observing the SO_2 and Sulfate Aerosol Plumes From the 2022 Hunga Eruption With the Infrared Atmospheric Sounding Interferometer (IASI)". In: *Geophysical Research Letters* 51(19). DOI: 10.1029/2023gl105565.

Simanjuntak, F. and T.-H. Lin (2022). "Monsoon Effects on Chlorophyll-a, Sea Surface Temperature, and Ekman Dynamics Variability along the Southern Coast of Lesser Sunda Islands and Its Relation to ENSO and IOD Based on Satellite Observations". In: *Remote Sensing* 14(7), page 1682. DOI: 10.3390/rs14071682.

Stenchikov, G. (2009). "The Role of Volcanic Activity in Climate and Global Change". In: *Climate Change*. Elsevier, pages 77–102. ISBN: 9780444533012. DOI: 10.1016/b978-0-444-53301-2.00004-x.

Stenchikov, G., A. Robock, V. Ramaswamy, M. D. Schwarzkopf, K. Hamilton, and S. Ramachandran (2002). "Arctic Oscillation response to the 1991 Mount Pinatubo eruption: Effects of volcanic aerosols and ozone depletion". In: *Journal of Geophysical Research: Atmospheres* 107(D24). DOI: 10.1029/2002jd002090.

Su, T., Z. Li, N. R. Henao, Q. Luan, and F. Yu (2024). "Constraining effects of aerosol-cloud interaction by accounting for coupling between cloud and land surface". In: *Science Advances* 10(21). DOI: 10.1126/sciadv.adl5044.

Timmreck, C., M. Toohey, D. Zanchettin, S. Brönnimann, E. Lundstadt, and R. Wilson (2021). "The unidentified volcanic eruption of ~1809: why it remains a climatic cold case". In: *EGU General Assembly 2021*. EGU21-5330. Online, 19–30 Apr 2021. DOI: 10.5194/egusphere-egu21-5330.

Toohey, M., Y. Jia, and S. Tegetmeier (2021). "Stratospheric residence time and the lifetime of volcanic aerosol". In: *EGU General Assembly 2021*. EGU21-12131. Online, 19–30 Apr 2021. DOI: 10.5194/egusphere-egu21-12131.

Trismidianto, D. Satiadi, W. Harjupa, I. Fathrio, Risyanto, E. Saufina, R. Muharsyah, D. E. Nuryanto, F. Nauval, D. F. Andarini, A. Puruwaningsih, T. Harjana, A. S. Praja, A. Witono, I. Juaeni, and B. Suhandi (2024). "Southerly Surge Impact on Rainfall Patterns in Southern Indonesia during Winter Monsoon and Madden–Julian Oscillation (MJO)". In: *Atmosphere* 15(7), page 840. DOI: 10.3390/atmos15070840.

Virtanen, A., J. Joutsensaari, H. Kokkola, D. G. Partridge, S. Blichner, Ø. Seland, E. Holopainen, E. Tovazzi, A. Lipponen, S. Mikkonen, A. Leskinen, A.-P. Hyvärinen, P. Zieger, R. Krejci, A. M. L. Ekman, I. Riipinen, J. Quaas, and S. Romakkaniemi (2025). "High sensitivity of cloud formation to aerosol changes". In: *Nature Geoscience* 18(4), pages 289–295. DOI: 10.1038/s41561-025-01662-y.

Whitty, R. C. W., E. Ilyinskaya, E. Mason, P. E. Wieser, E. J. Liu, A. Schmidt, T. Roberts, M. A. Pfeffer, B. Brooks, T. A. Mather, M. Edmonds, T. Elias, D. J. Schneider, C. Oppenheimer, A. Dybwad, P. A. Nadeau, and C. Kern (2020). "Spatial and Temporal Variations in SO_2 and PM2.5 Levels Around Kīlauea Volcano, Hawai'i During 2007–2018". In: *Frontiers in Earth Science* 8. DOI: 10.3389/feart.2020.00036.

Wirasatriya, A., R. D. Susanto, K. Kunarso, A. R. Jalil, F. Ramdani, and A. D. Puryajati (2021). "Northwest monsoon upwelling within the Indonesian seas". In: *International Journal of Remote Sensing* 42(14), pages 5433–5454. DOI: 10.1080/01431161.2021.1918790.

Wong, A., I. Roy, W. Yim, and A. Roy (2025). "Volcanic eruptions as a cause of extreme weather with special focus on Hong Kong". In: *Air Pollution and Related Health Risks*. Elsevier, pages 331–350. ISBN: 9780443239656. DOI: 10.1016/b978-0-443-23965-6.00012-5.

Wong, W. C. A., A. Roy, W. W.-S. Yim, and I. Roy (2023). "Can Volcanoes be Linked to Extreme Monsoon Rainfall in Asian Countries?" In: *Asia Oceania Geoscience Society (AOGS) 2023*. DOI: 10.13140/RG.2.2.26737.25446.

Wong, W. C. A. and W. W.-S. Yim (2022). *2021–2022 Tongan volcanic eruptions and record rainfall in eastern Australia and New Zealand*. Volcanoes Study Group, Hong Kong.

Yim, W. W.-S. (2005). "The El Chichón eruption and rainfall distribution in Hong Kong during 1982". In: *5th Asia–Pacific Space Geodynamics Meeting*.

Yu, P., R. W. Portmann, Y. Peng, C.-C. Liu, Y. Zhu, E. Asher, Z. Bai, Y. Lu, J. Bian, M. Mills, A. Schmidt, K. H. Rosenlof, and O. B. Toon (2023). "Radiative Forcing From the 2014–2022 Volcanic and Wildfire Injections". In: *Geophysical Research Letters* 50(13). DOI: 10.1029/2023gl103791.

Zheng, G., H. Su, M. O. Andreae, U. Pöschl, and Y. Cheng (2024). "Multiphase Buffering by Ammonia Sustains Sulfate Production in Atmospheric Aerosols". In: *AGU Advances* 5(4). DOI: 10.1029/2024av001238.